

AD-A249 993



ENTATION PAGE

Form Approved
OMB No. 0704-0188

2

estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the necessary data, reviewing the collection of information, Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 92-04-15		3. REPORT TYPE AND DATES COVERED Final 18 Jan 88 - 17 Jul 91	
4. TITLE AND SUBTITLE Dynamical Systems in Neuropharmacology				5. FUNDING NUMBER DAAL03-88-K-0018	
6. AUTHOR(S) Mark A. Geyer, Ph.D. Arnold J. Mandell, M.D.					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Regents of the University of California University of California, San Diego La Jolla, California 92093-0804				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709-2211				10. SPONSORING/MONITORING AGENCY REPORT NUMBER ARO 25701.12-LS	
11. SUPPLEMENTARY NOTES The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.				12b. DISTRIBUTION STATEMENT DTIC ELECTE MAY 04 1992 S D	
13. ABSTRACT (Maximum 200 words) Quantitative techniques found to be useful in characterizing the complex attractors of dynamical systems and the non-convergent distribution functions of statistical hierarchies were applied to the problem of drug-induced phase-transitions in patterns of exploratory behavior in the rat. Methods derived from ergodic theory and the application of scaling exponents proved to be particularly useful. These techniques provided discriminations among drug effects and insights into behavioral mechanisms that were not accessible using traditional psychopharmacological measures.					
14. SUBJECT TERMS Dynamical Systems, Ergodic Theory, Psychostimulants, Dopamine, Serotonin, Scaling Exponents				15. NUMBER OF PAGES 7	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL		

DYNAMICAL SYSTEMS IN NEUROPHARMACOLOGY

FINAL REPORT

MARK A. GEYER, Ph.D.

AND

ARNOLD J. MANDELL, M.D.

APRIL 15, 1992

U.S. ARMY RESEARCH OFFICE

DAAL03-88-K-0018

THE REGENTS OF THE UNIVERSITY OF CALIFORNIA

DEPARTMENT OF PSYCHIATRY

UNIVERSITY OF CALIFORNIA, SAN DIEGO

Accession For		
NTIS	CRA&I	<input checked="" type="checkbox"/>
DTIC	TAB	<input type="checkbox"/>
Unannounced		<input type="checkbox"/>
Justification		
By		
Distribution /		
Availability Codes		
Dist	Avail and/or Special	
A-1		



92-11865



02 4 29 131

DYNAMICAL SYSTEMS IN NEUROPHARMACOLOGY**FINAL REPORT**

MARK A. GEYER, Ph.D.
AND
ARNOLD J. MANDELL, M.D.

The original proposal for this project listed three principal goals:

1. Further elaboration and application of the quantitative techniques of modern dynamical systems theory and experimental mathematics to sequential data or "time series".
2. Numerical exploration of both smooth and discrete abstract dissipative dynamical systems as realistic low-dimensional compressions of the dynamical response pattern to increasing doses of stimulants.
3. Development of phase-space graphic computer programs for the qualitative description of changes in complexity induced by stimulant and other drugs in neuronal, neurochemical, and behavioral dynamical systems.

Data were to be generated in three laboratories studying the effects in rats of stimulants and other parametric perturbations on: (a) locus coeruleus, substantia nigra, and cortical neuron interspike intervals; (b) continuously HPLC(ec)-monitored levels of NE, MHPG, DA, DOPAC, HVA, 5HT, and 5HIAA in brainstem perfusion studies of freely moving rats; (c) digitally resolved temporal and spatial exploratory behavior in conjunction with (b). The laboratories to be involved were those of Drs. Stephen Foote (a), Ronald Kuczenski (b), and David Segal (c) of UCSD. The shared interests of these investigators in brainstem biogenic amine-mediated neurobiological mechanisms of behavior and the use of cocaine, amphetamine, and other pharmacological probes of mechanisms was expected to permit cross-level exploration for the dynamical universality predicted by the theory.

As the project progressed, some alterations in the emphasis of the project were required. The fundamental premise of the project was to assess the utility of applying non-linear dynamical methods to high-quality biological data. That general objective of the project was accomplished. However, the specific databases upon which these methods were proposed to be developed were not all used effectively. Specifically, the planned use of time series data derived from Dr. Kuczenski's HPLC analyses of neurochemical changes in freely moving rats never materialized. Similarly, no substantive progress was made using data from Dr. Segal's locomotor activity chambers. Some promising work has been accomplished in collaboration with Dr. Foote, using his measures of single unit responses of cells in the locus coeruleus, substantia nigra, and cortex of rats. This work is currently in manuscript form and should be submitted for publication within the year.

In some of our more strictly theoretical work, we demonstrated some dynamical consequences of making the formal characteristic of neuronal elements and their connectivities more realistic than those of modern neural computer algorithms [3]. Specifically, we studied two multiplicatively coupled logistic maps, each with the quadratic response characteristics of neurons with autoreceptor mechanisms, which iteratively exchange their outputs. The dynamic phenomenology of this two-neuron system, though noninvertible, demonstrated characteristics of both automorphisms of the real

line and diffeomorphisms of the plane, with smooth parametric transitions between them. We portrayed a unified parameter space containing the negative, positive, and mixed-coupled regimes, categorized the phase space structures using characteristic exponent inequalities, explored representative parametric paths, demonstrated generic bifurcation sequences, and reported number theoretic ordering in parameter space. We conjectured that most, if not all, "middle-layer-like" brain systems, in contrast to primary sensory and motor information transport systems, are dominated by intrinsic dynamics of the sort we demonstrated. According to this view, external input serves as a perturbation of these already ongoing complex systems. The intrinsic instabilities of these middle-layer dynamics return the unpredictability required by theoretical studies of real brain functions to models of neural networks. We demonstrated a variety of parameter-dependent intrinsically ordered attractor sequences and number theoretic series which emerge from the global dynamics and exemplify the rich dynamical "machine" language potential for specific coding by brain-like information transport and learning algorithms.

Another theoretical development was pursued to examine the statistical properties of real macromolecular sequences using an approximation based on the composition of two ergodic sets [2]. Both varied and reliable mechanisms of information transport in molecular biological systems can be obtained from the universality properties of continuous and discrete dynamical systems whose output has the general properties of a set which combines sensitivity to initial conditions and indecomposability with a positive density of periodic points. This approach required a revision of information theory for macromolecular systems dictated by their failure to fulfill the conditions of Shannon's existence proof: there exists a one-to-one mapping that makes the channel capacity finite. Relaxation studies of polypeptides, proteins, and nucleotides demonstrate a lack of characteristic times. Using calcitonin as the exemplar, a more appropriate statistical approach to macromolecular coding was explored using a hyperbolic distribution with an infinite multiplicity of moments, multiplicative noise, and belonging to the family of convolutionally stable distributions without finite mean or variance [2]. This work used a phase space reconstruction technique to relate the potencies of important calcitonin analogues to the global "tightness" of the vorticity.

The most significant work related to concrete biological systems was based on the data from Dr. Geyer's Behavioral Pattern Monitor (BPM) system, which provided the pilot data on behavioral measures presented in the original proposal. The extensive database derived from the BPM system proved to be ideal for addressing the conceptual questions originally proposed. Thus, we had considerable success by carefully examining data from studies of the behavioral profiles of psychoactive drugs as characterized in the BPM system. In particular, we developed new methods based on procedures developed in the context of theoretical physics and non-linear dynamics. Using a k-D splitting algorithm, we described the initial demonstrations of the utility of complexity and entropy measures in biological applications, using data from the BPM system [7]. Measures of complexity derived from ergodic theory of dynamical systems were developed and applied to an exemplary data set describing locomotor movements of rats in a bounded space. A symbolic dynamical system was obtained by partitioning the event space into equally probable partition elements, using a k-dimensional tree. The measures calculated from the symbolic sequences included the topological entropy (ht)--i.e., the rate of increase of all possible sequences with increasing sequence length--and the metric entropy (hm)--i.e., the rate of increase of all likely sequences with increasing sequence length. These measures were used to assess changes in rat locomotor behavior as recorded in the behavioral pattern monitor (BPM) that were induced by amphetamine (0.25, 0.50, 1.0, or 2.0 mg/kg) and 3,4-methylenedioxymethamphetamine (MDMA; 1.25, 2.5, 5.0, or 10.0 mg/kg). Amphetamine increased the mean activity, ht, and hm. MDMA resulted in a monotonic dose-response curve for

activity but exhibited a biphasic dose response in ht and hm. In particular, some animals in the higher dose groups showed a ht in the range of the saline controls, whereas other animals exhibited a significantly reduced ht and a greater decrease in hm, suggesting that two different behavioral reactions coexist within the same higher dose range of MDMA. An important implication of our method is that, in applied ergodic measure-theoretic approaches, the partition that determines the elements of the symbolic dynamical system should not be specified a priori on abstract mathematical grounds but should be chosen relative to its significance with respect to the data set in question. Here, the animal constructs its own spatiotemporal partition in behavioral phase space.

In addition to the complexity and entropy measures, we also made great progress in the use of scaling measures to provide more sensitive characterizations of drug effects. We used scaling measures to provide more sensitive characterizations of drug effects [4,6]. Conceptually, these measures are derived from the concept of dimensionality in the field of fractal geometry. We have defined two global scaling measures from our BPM data, one characterizing the temporal structure of the animal's behavior, the other being sensitive to changes in the geometrical structure of the animal's locomotor path. These two measures enable us to assess quantitatively similarities and differences between different drug treatments. The first full description of this approach appeared in Psychopharmacology [6]. This paper details the derivations and applications of the temporal scaling exponent, α , and the spatial scaling exponent, d , and introduces the d - α plane. Basically, α assesses the level of activity of an animal and has proven to be more consistent and sensitive than the traditional measure of photobeam breaks, presumably because it utilizes all the available information. The spatial scaling exponent d describes the roughness of the path traversed by the animal - it assesses the average geometrical information contained in locomotor activity in a manner that is independent of changes in the amount of activity. The d - α plane provides a condensed way to present and visualize a large amount of information about the nature of a drug's effects across a range of doses and facilitates comparisons between the effects of different drugs. To illustrate this feature of our approach, we compared the effects of a variety of psychostimulant drugs and used these comparisons to address hypotheses regarding the behavioral processes affected by these drugs [4,5,6].

A variety of psychoactive substances (amphetamine, nicotine, scopolamine, apomorphine, lisuride, and MDMA) were tested to examine whether a scaling hypothesis is appropriate for the description of the amount and the structure of rat locomotor paths recorded in the BPM [6]. The analytical approach was based on the assumption that the scaling behavior of a few collective variables may characterize sufficiently changes in the animal's behavior induced by different drugs. The temporal scaling exponent α , describing the ratio of fast to slow responses in the BPM, sensitively reflected the different stimulant properties of the substances. The spatial scaling exponent d , which relates the average path length to the resolution used to measure consecutive responses, was found to discriminate substances that had been separated previously via qualitative descriptions. Several behavioral response categories emerged from comparisons of the locations of different drugs on a two-dimensional d - α plane. Scopolamine, MDMA, lisuride, and high doses of apomorphine increased α while decreasing d , whereas amphetamine, nicotine, and caffeine produced an increased α with no change or an increase in d . Stereotypes could be identified on the opposite ends of the spatial scaling exponent scale and were interpreted as reflecting two kinds of perseveration. These results suggest that scaling approaches can be used to assess quantitatively the state of the animal based on its locomotor behavior and that the exponents can serve as collective variables providing a macroscopic description based on the microscopic elements of behavior.

Using both our traditional measures and the newer theoretically driven measures, we explored the mechanisms contributing to the locomotor hyperactivity induced by MDMA [1,5]. We

discovered that the profound locomotor activation produced by MDMA and related drugs is not due primarily to dopamine release, but is due to serotonin release [1]. We blocked MDMA-induced hyperactivity with the selective serotonin uptake blockers fluoxetine, sertraline, or zimelidine or with the serotonin synthesis inhibitor PCPA. By contrast, these pretreatments either potentiated or had no effect on the hyperactivity induced by the dopamine releasing agent amphetamine. Conversely, the catecholamine synthesis inhibitor AMPT, which blocks the effects of amphetamine, did not affect the response to MDMA. We applied our scaling and complexity measures to this problem [5]. The use of scaling measures enabled us to further differentiate the effects of MDMA from those of either amphetamine or hallucinogens. In addition, comparisons of the behavioral effects of MDMA across both repeated tests and repeated administrations of the drug confirmed that our spatial scaling exponent, d , was affected differently from more traditional measures. Furthermore, d proved to be the most sensitive of our various measures of the effects of serotonin releasers on the spatial structure of the locomotor paths exhibited by the animals. The combination of our new and old measures of drug effects on behavior clearly provides more information than can be obtained by reliance on any one set of measures.

The d - α plane was also used to systematically compare various methylenedioxy-substituted phenethylamines having varying potencies as releasers of dopamine or serotonin [5]. This analytical approach enabled us to make discriminations between drugs and specific enantiomers of drugs that have previously required much more elaborate and time-consuming procedures such as drug-discrimination tests. With this approach, we can characterize and quantify different drug effects in a single test session using a procedure that does not require repeated administrations of the drug to the animals. Thus, the effects of chronic versus acute exposures to the drugs can now be assessed reliably.

In further work, we developed modified versions of both the scaling measures and the previously published entropy measures. These extensions include the generalization of the entropy as well as the spatial scaling exponent, $S(h)$ and $f(d)$ respectively. These extensions allow us to not only describe the average scaling measure but enables us to assess quantitatively the contribution of different subsets to the overall behavior. This approach is based on the idea that the animals engage in different "types" of behavior which can be quantitatively described by different local scaling exponents. For example, a rat moving along the wall in a straight path may also engage in highly repetitive head movements or gnawing stereotypies. We have observed precisely this type of behavior in rats treated with 20.0 mg/kg cocaine. However, the averaging procedure yielding the average spatial scaling exponent d would not capture the distinctly different characteristics of these behavioral "subsets". Thus, using the spectrum of spatial scaling exponents, $f(d)$, we can now assess these behavioral subsets in addition to the average geometrical path structure [4]. The singularity spectrum of local entropies ($S \sim (\sim \alpha \sim)$) computed from the microcanonical dynamical partition function was used to describe the sequential dynamical structure of rat locomotor paths under the influence of cocaine. The approach used was analogous to the microcanonical ensemble of statistical mechanics and permitted the sensitive detection of "phase transition" like behavior. The results suggest that the behavioral sequences exhibited by the rats are determined by two processes, each with its own entropy ($\sim \alpha_{\{l\}} \sim, \sim \alpha_{\{h\}} \sim$). These processes are changed simultaneously by cocaine, suggesting that they coexist as local repellers with overlapping insets and undergo a change from second to first order phase transition [8]. With this approach we may be able to differentiate between different drugs, e.g. MDMA, high doses of cocaine, or scopolamine, generating an overall increase in straight path patterns but acting presumably via very different mechanisms. These elaborations have already led to even more precise insights into mechanisms underlying the

behavioral effects of psychoactive drugs.

PUBLICATIONS SUPPORTED IN PART BY THE AWARD

- Mandell, A.J.: An unstable singularity theory of molecular biological coding: Calcitonin's structures and potencies. In: *Dynamic Patterns in Complex Systems*. Eds. J.A.S. Kelso, A.J. Mandell, and M.F. Shlesinger. World Scientific, Singapore, pp 219-235, 1988.
- Paulus, M.P., Gass, S.F. and Mandell, A.J.: A realistic, minimal "middle-layer" for neural networks. *Physica D*, 40:135-155, 1989.
- Paulus, M.P. and Geyer, M.A.: A temporal and spatial scaling hypothesis for the behavioral effects of psychostimulants. *Psychopharmacology*, 104(1):6-16, 1991.
- Paulus, M.P., Geyer, M.A. and Mandell, A.J.: Statistical mechanics of a neurobiological dynamical system: The spectrum of local entropies ($S(\alpha)$) applied to cocaine-perturbed behavior. *Physica A*, 174:567-577 1991.
- Paulus, M.P. and Geyer, M.A.: A scaling approach to find order parameters quantifying the effects of dopaminergic agents on unconditioned motor activity in rats. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 15(6):903-919, 1991.
- Paulus, M.P. and Geyer, M.A.: The effects of MDMA and other methylenedioxy substituted phenylalkylamines on the structure of rat locomotor activity. *Neuropsychopharmacology*, in press.
- Paulus MP; Geyer MA; Gold LH; Mandell AJ. Application of entropy measures derived from the ergodic theory of dynamical systems to rat locomotor behavior. *Proceedings of the National Academy of Sciences of the United States of America*, 1990 Jan, 87(2):723-7.

PERSONNEL

The following personnel participated in the project: Mark A. Geyer, Ph.D., Suzanne Knapp, Ph.D., Arnold J. Mandell, M.D., Steve Gass, and Martin Paulus, M.D.

BIBLIOGRAPHY

- Callaway, C.W., Nichols, D.E., Paulus, M.P. and Geyer, M.A.: Serotonin release is responsible for the locomotor hyperactivity in rats induced by derivatives of amphetamine related to MDMA. In: *Serotonin: Molecular Biology, Receptors and Functional Effects*, (J.R. Fozard & P.R. Saxena, eds), Birkhauser, Basel, 491-505, 1991.
- Mandell, A.J.: An unstable singularity theory of molecular biological coding: Calcitonin's structures and potencies. In: *Dynamic Patterns in Complex Systems*. Eds. J.A.S. Kelso, A.J. Mandell, and M.F. Shlesinger. World Scientific, Singapore, pp 219-235, 1988.
- Paulus, M.P., Gass, S.F. and Mandell, A.J.: A realistic, minimal "middle-layer" for neural networks. *Physica D*, 40:135-155, 1989.
- Paulus, M.P. and Geyer, M.A.: A scaling approach to find order parameters quantifying the effects of dopaminergic agents on unconditioned motor activity in rats. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 15(6):903-919, 1991.
- Paulus, M.P. and Geyer, M.A.: The effects of MDMA and other methylenedioxy substituted phenylalkylamines on the structure of rat locomotor activity. *Neuropsychopharmacology*, in press.
- Paulus, M.P. and Geyer, M.A.: A temporal and spatial scaling hypothesis for the behavioral effects

of psychostimulants. *Psychopharmacology*, 104(1):6-16, 1991.

Paulus MP; Geyer MA; Gold LH; Mandell AJ. Application of entropy measures derived from the ergodic theory of dynamical systems to rat locomotor behavior. *Proceedings of the National Academy of Sciences of the United States of America*, 1990 Jan, 87(2):723-7.

Paulus, M.P., Geyer, M.A. and Mandell, A.J.: Statistical mechanics of a neurobiological dynamical system: The spectrum of local entropies ($S(\alpha)$) applied to cocaine-perturbed behavior. *Physica A*, 174:567-577 1991.